


Crystal Structure of the Sodium Cobaltate Deuterate Superconductor $\text{Na}_x\text{CoO}_2 \cdot 4x\text{D}_2\text{O}$ ($x \approx 1/3$)

J. D. Jorgensen, M. Avdeev, D. G. Hinks, J. C. Burley, S. Short, Materials Science Division, ANL

Motivation: Determine the crystal structure of the newly-discovered sodium cobaltate deuterate superconductor $\text{Na}_x\text{CoO}_2 \cdot 4x\text{D}_2\text{O}$ ($x \approx 1/3$). In particular, understand how water is incorporated and what this dictates about the chemistry of the superconducting compound.

Results:


- A single-phase deuterated sample was made by first determining the equilibrium vapor pressure for the deuterated superconducting phase.
- Neutron diffraction data were collected on the SEPD at IPNS.
- Fourier mapping revealed the general location of D_2O molecules.
- Rietveld refinement provided a detailed average structural model.
- Chemical bonding arguments led to an ordered model that explains additional scattering seen in the neutron diffraction data.



Impact: This result shows that the new superconductor is specific hydrated compound with four fully hydrogen-bonded water molecules coordinated to each Na ion, not an intercalated compound with independent and continuously variable Na and water contents or a compound with "ice" layers between the Na and CoO_6 layers. Different hydrated structures (e.g., with 6-coordinated Na) could form at other Na and water contents.

Future Work: Identify and explore the properties of other hydrated compounds in this family. Search for other layered compounds that are metallic, and superconducting, when hydrated.

Crystal Structure of the Sodium Cobaltate Deuterate Superconductor $\text{Na}_x\text{CoO}_2 \cdot 4x\text{D}_2\text{O}$ ($x \approx 1/3$),
J. D. Jorgensen, M. Avdeev, D. G. Hinks, J. C. Burley, S. Short, cond-mat/0307627



BES - DOE

This work was supported by the U. S. Department of Energy, Basic Energy Sciences, under contract W-31-109-ENG-38.

MSD - ANL

